

Intrauterine insemination

patient information

INTRODUCTION	3
THE MENSTRUAL CYCLE	4
WHAT IS INSEMINATION?	5
INITIATING INSEMINATION TREATMENT	6
INSEMINATION TREATMENT PROCEDURE	7
Step 1: Monitoring the cycle	
Step 2: Insemination	
Step 3: Pregnancy following insemination treatment	
PSYCHOLOGICAL SUPPORT	12
RELAXATION THERAPY	13
CONTACT DETAILS	13

This brochure provides information on intrauterine insemination (IUI).


Should you have any further questions after reading this brochure, please do not hesitate to contact us. The doctors and midwives at the Leuven University Fertility Centre will be happy to provide further information.

THE MENSTRUAL CYCLE

The start of the menstrual cycle coincides with the presence of several small **follicles**. One of these follicles will grow and mature within the ovary over the course of the first 14 days.

As the follicle grows it will produce more and more hormones (oestrogen). The high oestrogen concentration will affect the endometrial lining and cervix. **Ovulation** will follow under the influence of the luteinising hormone. The mature follicle will tear and release the egg, approximately 14 days before the end of the menstrual cycle.

When the egg is released it is captured in the oviductal ampulla. The egg proceeds through the fallopian tube **as a result of the rhythmic contractions of the uterus and fallopian tube**.



What remains of the follicle after ovulation develops into a corpus luteum under the influence of the luteinising hormone. The corpus luteum secretes progesterone, a hormone that maintains the endometrium. If the egg is not fertilised after ovulation the corpus luteum will not survive. Progesterone production will decrease and the endometrial lining will break down, resulting in menstruation.

Insemination does not necessarily coincide with a normal menstrual cycle:

- ✓ Hormonal medication can be used to stimulate a cycle.
- ✓ The sperm sample is capacitated and the concentration and motility checked in the laboratory.
- ✓ The processed sperm sample is injected directly into the uterine cavity around the time of ovulation.

WHAT IS INSEMINATION?

Intrauterine insemination (IUI) is a type of fertility treatment in which processed sperm cells are inserted into the uterus at the time of ovulation.

INITIATING INSEMINATION TREATMENT

You can contact the Leuven University Fertility Centre at your own initiative or with a referral from a doctor.

Your initial consultation with the fertility centre will provide room for a [detailed consultation](#). In preparation of this initial consultation, you will first have to complete a [medical questionnaire](#) at home. The fertility consultant will ask specific questions to get a clear understanding of your situation. Sometimes this consultation will be prepared by a specialist midwife. You will then be given information on [additional examinations](#) involving you and your partner. The examination phase may take up to two to three months.

Once all examinations have been completed [a consultation](#) will be arranged with the fertility doctor to discuss the results of the examinations. Insemination treatment can be initiated providing at least one fallopian tube is unobstructed and the sperm sample is of acceptable quality.

Before starting fertility treatment, you will receive information on the various stages of the treatment. Information videos provide more explanation prior to your consultation with the [midwife](#). This is also when the [signed contracts](#) are handed over to schedule the start of the treatment.



INSEMINATION TREATMENT PROCEDURE

STEP 1: MONITORING THE CYCLE

Insemination treatment can be monitored during a natural cycle. The menstrual cycle is monitored by means of ultrasound follicle measurements and blood samples until the time of ovulation.

The ovaries may also be stimulated using hormone medication.

The following is a summary of the different medications:

1. Clomiphene citrate (Clomid[®], Letrozole[®])

Clomiphene citrate is an anti-oestrogen that stimulates the release of the follicle stimulating hormone, which is vital for **follicle growth**. The treatment consists in taking tablets **for five days at a specific stage of the cycle**.

Follicle growth is monitored by means of several blood samples and ultrasound follicle measurements until the follicle has matured.



The start of the treatment is always discussed with your doctor at the consultation.

2. Gonadotropins (Menopur[®], Gonal F[®])

If a natural cycle or treatment with hormone tablets does not result in pregnancy, hormone injections can be administered before insemination treatment.

Gonadotropins contain follicle stimulating hormone (FSH), with or without luteinising hormone (LH), which **stimulates the growth of follicles**. The hormones are administered **daily via subcutaneous injections**, which you, your GP or the home nurse can perform. Follicle growth will be monitored by means of several blood samples and ultrasound follicle measurements.

Gonadotropins are administered during maximum six treatments. In most cases a rest cycle without medication is introduced after one or two treatment cycles.



The Ministerial Decree (MD) dated 14 September 2006 regulates the reimbursement of gonadotropins within the framework of controlled ovarian stimulation with or without insemination. The MD stipulates that gonadotropins will be refunded subject to specific indications. Your doctor will be able to provide further information and the necessary documentation.


STEP 2: INSEMINATION

Synthetic chorionic gonadotropin (Ovitrelle®) promotes ovulation. Ovitrelle® is administered via a subcutaneous injection into the abdomen. The insemination will be scheduled around the time of ovulation.

On the day of the insemination your partner will have to deliver a sperm sample to the laboratory, where it will be processed.


If a frozen or donor sample is used it will be thawed on the day of the insemination.

The processed sperm is then injected into the uterus via a thin catheter. The procedure is not painful. You will be able to return to your daily activities after the insemination. If the insemination is in a weekend, supplements may be charged.



Risks

The use of hormone medication can promote the growth of several follicles. This is referred to as overstimulation, which is associated with an increased likelihood of multiple pregnancy if more than two follicles mature. In such cases there will be no insemination and you will have to wait until the next cycle. Careful monitoring and a timely reduction in the medication can limit the risk of overstimulation.



Ovary with one follicle



Ovary with several follicles

STEP 3: PREGNANCY FOLLOWING INSEMINATION TREATMENT

The likelihood of pregnancy following insemination treatment is 16% on average. The use of gonadotropins increases the likelihood of pregnancy.

The risk of miscarriage is between 10 and 15% for women aged between 30 and 34. For women aged between 35 and 39 this risk increases to 18 to 25%.

And finally there is a small risk of ectopic pregnancy of 3%.

PSYCHOLOGICAL SUPPORT

Fertility problems or the start of fertility treatment are extremely emotional experiences. You may be subject to various emotions and thoughts, continually changing in intensity and direction, at different times. Often, talking to your partner or someone close to you may help. If you feel there are still a number of unanswered questions you can request a consultation with a specialized psychologist.

Our psychologist will be happy to talk to you individually, or together with your partner, covering all the different aspects of fertility problems. Ample time will be taken to consider your specific concerns and the individual and relational decision-making process in detail. How can you talk about your experiences with your partner or your nearest and dearest? How do you manage insecurities, the highs and lows of the treatment process? Sometimes it may be a relief to be told that your feelings and thoughts are a normal part of the journey. Counseling is always possible, during the examination stage, the actual treatment or after its termination.

RELAXATION THERAPY

If you are under a lot of stress as a result of the treatment or a combination of different factors, our psychomotor therapist will be happy to provide support.

Psychomotor therapy is a type of therapy which **focuses on the body and body awareness**. This may help when stress manifests itself in physical complaints such as muscle tension, anxiety, sleeping badly, no longer being able to relax, pain, and hyperventilation. Your specific symptoms are examined and the results used to jointly find a meaningful approach to deal with your particular situation. Various tips and techniques will be provided, tailored to your specific requirements, to help you along the way.

CONTACT DETAILS

Leuven University Fertility Centre

Medical queries: 016 34 36 24

Administrative queries: 016 34 36 50

www.uzleuven.be/lufc

Psychologist: Anneleen Rasquin

Physiotherapist: Gillian Demin

© May 2025 UZ Leuven

This text and these illustrations can only be copied subject to prior authorisation from the UZ Leuven communications department.

Design and implementation

This text was written by the Leuven University Fertility Centre in cooperation with the communications department.

This brochure is also available at www.uzleuven.be/en/brochure/700999.

Please send comments or suggestions relating to this brochure to communicatie@uzleuven.be.

Publisher
UZ Leuven
Herestraat 49
3000 Leuven
tel. 016 33 22 11
www.uzleuven.be

 mynexuzhealth



Consult your medical record
via nexushealth.com
or download the app

